

L4 ANSWER 1 OF 1 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 AN 2001:322750 BIOSIS  
 DN PREV200100322750  
 TI Generation of CD8+ T cell lines with specific cytotoxicity for autologous Chronic Lymphocytic Leukemia B cells.  
 AU Deforce, Dieter L. (1); Chu, Peter (1); Rassenti, Laura Z. (1); Mendoza, Robert (1); Raz, Eyal (1); Kipps, Thomas J. (1)  
 CS (1) Hematology/Oncology, UCSD School of Medicine, La Jolla, CA USA  
 SO Blood, (November 16, 2000) Vol. 96, No. 11 Part 1, pp. 368a. print.  
 Meeting Info.: 42nd Annual Meeting of the American Society of Hematology San Francisco, California, USA December 01-05, 2000 American Society of Hematology  
 . ISSN: 0006-4971.  
 DT Conference  
 LA English  
 SL English  
 AB Although a lot of effort has been put in the expansion of autologous CD8+ T cells in response to CD40L activated Chronic Lymphocytic Leukemia (CLL)-B cells both by our group and by others (Blood 93:1992, 1999), the responding T cells are mainly CD4+. Experiments with normal donors show that normal CD8+ T cells can proliferate in response to CD40L activated CLLB (CD40L-CLLB) cells. Both our group (unpublished data) and others (Blood 95:999, 2000) have shown that CpG-motif containing phosphorothioate oligonucleotides (CpG-ODN) activate CLLB cells to express co-stimulatory molecules and that it is synergistically with CD40L activation. CpG-ODN also activates dendritic cells. Based upon this we investigated if CpG-ODN could promote long term proliferation of autologous CD8+ T cells in response to CD40L-CLLB cells. For this purpose PBMC's of CLLB patients were cultured with or without CpG-ODN (TGACTGTGAACGTTTCGAGATGA) or non-CpG-ODN (TGACTGTGAAGTTTAGAGATGA or TGACTGTGAATGTTTAGAGATGA). Initial anergy of the T cells was overcome by a onetime stimulation with immobilized mAb's to CD3 (OKT3) and CD28 (TN228), for subsequent stimulation CD40L-CLLB cells were added every 3 days. The cultures were followed over at least 4 weeks, including their T cell receptor (TCR) repertoire. By week 3 the cultures with CpG-ODN showed on average (n=5) a double amount of CD8+ T cells compared to the controls or the cultures with the non-CpG-ODN. The T cell cultures got more restricted to certain Vbeta TCR, showing that they got more oligoclonal. The involvement of dendritic cells was shown in a depletion experiment: the effect of CpG-ODN could be completely abolished by removing all adherent cells. The cultures in the presence of adherent cells and CpG-ODN showed about a 3 fold increase in the production of IFNgamma and IL-6 compared to either CpG-ODN or co-culture with adherent cells alone. With CpG-ODN we were able to show CD1a+ cells in CLL-PBMC's after 4 days in culture, whereas none were detected in the absence of CpG-ODN. These CD1a+ dendritic cells were phenotyped by FACS (CD40+, CD86+, CD4+, CD54 +++, HLA-DR+++). Cytotoxicity (CTL) assays with the purified CD8+ T cells showed that these cells caused about 50% of the autologous CD40L-CLLB cells to undergo apoptosis at a effector to target ratio (E/T) of 10:1. This killing could be inhibited by

W6/32, a mAb specific for a non-polymorphic epitope of HLA class I, and  
by mAb's specific for the Vbeta TCR which was increasing in response to  
co-culture with autologous CD40L-CLLB cells. Leading to the conclusion  
that CpG-ODN is capable of stimulating long term proliferation of CD8+ T  
cells with high potency to kill autologous CLLB cells via a TCR  
restricted  
manner.